

多穗金粟兰中一个新的酚苷^{*}

冉新辉^{1,2}, 倪伟², 魏刚³, 陈昌祥², 刘海洋^{2**}

(1 广州中医药大学第一临床医学院, 广东 广州 510405; 2 中国科学院昆明植物研究所植物化学与西部植物资源持续利用国家重点实验室, 云南 昆明 650204; 3 广州中医药大学新药研究中心, 广东 广州 510006)

摘要: 从多穗金粟兰 (*Chloranthus multistachys*) 全株的甲醇提取物中分离得到 1 个新的酚苷, 通过波谱技术鉴定其结构为丁香酸-4-*O*- α -L-鼠李吡喃糖苷 (**1**)。同时还首次从该植物中分离得到两个已知倍半萜内酯和一个已知木脂素: 1 β , 4 β -二羟基-5 α , 8 β -二氢-7 (11) *Z*-桉叶烯-8, 12-内酯 (**2**), 1 β , 4 α -二羟基-5 α , 8 β -二氢-7 (11) *Z*-桉叶烯-8, 12-内酯 (**3**), (–)-(7*S*, 8*R*)-dihydrodehydrodiconiferyl alcohol (**4**)。

关键词: 多穗金粟兰; 金粟兰科; 酚苷; 倍半萜内酯

中图分类号: Q 946

文献标识码: A

文章编号: 0253-2700(2010) 01-083-04

A New Phenolic Glycoside from *Chloranthus multistachys* (Chloranthaceae)

RAN Xin-Hui^{1,2}, NI Wei², WEI Gang³, CHEN Chang-Xiang², LIU Hai-Yang²

(1 The First Affiliated Hospital of Guangzhou University of Traditional Chinese Medicine, Guangzhou 510405, China;

2 State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany,

Chinese Academy of Sciences, Kunming 650204, China; 3 Research Center of New Drug,

Guangzhou University of Traditional Chinese Medicine, Guangzhou 510006, China)

Abstract: A new phenolic glycoside, named syringic acid-4-*O*- α -L-rhamnopyranoside (**1**), together with three known compounds were isolated from the MeOH extract of the whole plants of *Chloranthus multistachys*. Their structures were elucidated on the basis of extensive spectroscopic. To the best of our knowledge, the three known compounds, 1 β , 4 β -dihydroxy-5 α , 8 β (*H*)-eudesm-7 (11) *Z*-en-8, 12-olide (**2**), 1 β , 4 α -dihydroxy-5 α , 8 β (*H*)-eudesm-7 (11) *Z*-en-8, 12-olide (**3**), and (–)-(7*S*, 8*R*)-dihydrodehydrodiconiferyl alcohol (**4**) were all isolated from this plant for the first time.

Key words: *Chloranthus multistachys*; Chloranthaceae; Phenolic glycoside; Sesquiterpene lactones

The plants of genus *Chloranthus* (Chloranthaceae), comprising 15 species, are mainly distributed in the east of Asia (Zhou, 1993). Most of these plants have been using as important medicinal herbs for the treatment of tumors, rheumatisms, and emmeniopathy in traditional Chinese medicine (Jiangsu New Medical College, 1977). Previous

phytochemical investigation on this genus revealed the presence of monoterpenoids, sesquiterpenoids, bisesquiterpenoids, diterpenoids, ligans, and flavone glycosides (Wu *et al.*, 2007; Yang and Yue, 2006; Yang *et al.*, 2007; 2008; Wang *et al.*, 2008; Xu *et al.*, 2007; Li *et al.*, 2008). *Chloranthus multistachys* is a perennial herb

* Foundation items: This work was financially supported by the Young Academic and Technical Leader Raising Foundation of Yunnan Province (No. 2008PY066)

** Author for correspondence; E-mail: haiyangliu@mail.kib.ac.cn; Tel: +86-871-5223246

Received date: 2009-10-12, Accepted date: 2009-11-23

作者简介: 冉新辉 (1984—) 女, 在读硕士研究生, 主要从事天然药物化学研究。

which mainly distributes over the south of China and its roots has been used to treat bone fracture in China as folk medicine (Editorial Committee of the Administration Bureau of Traditional Chinese Medicine, 1998). One sesquiterpenoid, three bisesquiterpenoids, and nine *ent*-kaurane diterpenoids have been isolated from this plant (Yang and Yue, 2006; 2008). As a part of our systematic research works on the genus (Teng *et al.*, 2009), the chemical constituents of *C. multistachys* were investigated and a new

phenolic glycoside, syringic acid-4-*O*- α -L-rhamnopyranoside (**1**) was obtained, along with three known compounds, 1 β , 4 β -dihydroxy-5 α , 8 β (*H*)-eudesm-7 (11) *Z*-en-8, 12-olide (**2**) (Yang *et al.*, 2007), 1 β , 4 α -dihydroxy-5 α , 8 β (*H*)-eudesm-7 (11) *Z*-en-8, 12-olide (**3**) (Yang *et al.*, 2007), and (–)-(7*S*, 8*R*)-dihydrodehydrodiconiferyl alcohol (**4**) (Zhu *et al.*, 2008) (Fig. 1). In this paper, we report the isolation and structural elucidation of the new phenolic glycoside.




Fig. 1 Structures of compounds **1-4**

Results and Discussion

Compound **1** was isolated as an amorphous powder and showed the molecular formula $C_{15}H_{19}O_9$, with 6 degrees of unsaturation, by the negative HR-ESI-MS at m/z 343.1040 $[M-H]^-$. The IR spectrum displayed peaks indicating hydroxy groups (3422 cm^{-1}), benzene rings ($1592, 1502, 1460\text{ cm}^{-1}$), and carboxyl groups (1700 cm^{-1}). UV absorptions at λ_{\max} 211, 254 nm suggested that **1** had an excellent conjugated system. Analysis of the ^{13}C -NMR and DEPT spectra (Table 1) of **1** indicated one carbonyl (δ_{C} 169.6, s), two methoxys (δ_{C} 56.5, q), two sp^2 methines (δ_{C} 107.7, d), and four sp^2 quaternary carbons [δ_{C} 128.0 (s), 154.4 ($\times 2$, s), 139.7 (s)], which showed the presence of syringic acid moiety in this compound (Shao *et al.*, 2008). The signals at δ_{H} 5.33 (d, H-1'), 4.14 (m, H-2'), 3.89 (m, H-3'), 3.44 (m, H-4'), 4.25 (m, H-5'), 1.20 (d, H-6') in the ^1H -


NMR spectrum and signals at δ_{C} 103.3 (d, C-1'), 71.9 (d, C-2'), 72.1 (d, C-3'), 73.6 (d, C-4'), 71.2 (d, C-5'), 17.8 (q, C-6') in the ^{13}C -NMR spectrum (Table 1) of **1** showed a L-rhamnose in this compound. The anomeric configuration of the rhamnopyranosyl was determined as α -oriented on the ground the chemical shift values of the C-3' (δ_{C} 72.1) and C-5' (δ_{C} 71.2) to those of the corresponding carbons of methyl α - and β -rhamnopyranoside (Kasai *et al.*, 1979). The L-rhamnose was linked to C-4 by the HMBC correlation from H-1' to C-4 as shown in Fig. 2. Based on above mentioned evidence, the structure of **1** was identified to be syringic acid-4-*O*- α -L-rhamnopyranoside.

Experimental

General experimental procedures Optical rotation were taken on a Horiba SEAP-300 polarimeter. UV spectra were obtained on a Shimadzu UV-2401PC spectrophotometer. IR spectra were measured with a Bio-Rad FTS-

Table 1 ^1H and ^{13}C NMR data of compound **1** in CD_3OD (400 MHz and 100 MHz, δ in ppm)

No.	δ_{H} (Multiplicity, J in Hz)	δ_{C}	HMBC (H \rightarrow C)
1		128.0 (s)	
2, 6	7.33 (2H, s)	107.7 (d)	C-1, 3, 7, 4
3, 5		154.4 (s)	
4		139.7 (s)	
7		169.6 (s)	
1'	5.33 (1H, d, 1.7)	103.3 (d)	C-2', 4'
2'	4.14 (1H, dd, 1.7, 3.3)	71.9 (d)	C-4', 1'
3'	3.89 (1H, dd, 3.3, 9.6)	72.1 (d)	C-4'
4'	3.44 (1H, t, 9.6)	73.6 (d)	C-3', 6'
5'	4.25 (1H, m)	71.2 (d)	C-1', 4', 6'
6'	1.20 (3H, d, 6.3)	17.8 (q)	C-4', 5'
3, 5-OMe	3.87 (6H, s)	56.5 (q)	C-3, 5

Fig. 2 Key HMBC Correlations of compound **1**

135 spectrometer with KBr pellets. FAB mass spectra were obtained on a VG Auto spec-3000 spectrometer and high-resolution ESI mass spectra were recorded on an API Qstar Pulsar LC/TOF instrument. NMR spectra were measured in CD_3OD and recorded on a Bruker AM-400 or a DRX-500 NMR spectrometer with TMS as an internal standard. Separation and purification were performed by column chromatography on silica gel (200–300 mesh, Qingdao Marine Chemical Inc., Qingdao, People's Republic of China), RP-18 (Merck), and Sephadex LH-20 (Amersham Biosciences, Sweden). Fractions were monitored by TLC, and spots were visualized by heating silica gel plates sprayed with 10% H_2SO_4 in EtOH.

Plant material The whole plants *Chloranthus multistachys* was collected in August 2007 from Xinning, Hunan Province, P. R. China, and identified by Dr. En-De Liu of Kunming Institute of Botany. A voucher specimen (No. HY0001) was deposited at the State Key Laboratory of Phy-

tochemistry and Plant Resources in West China.

Extraction and isolation Dried and powered whole plants of *C. multistachys* (20 kg) were extracted three times with CH_3OH under reflux. The filtrate was evaporated under reduced pressure to give a residue, which was suspended into water and partitioned with EtOAc. The EtOAc portion (1300 g) was subjected to column chromatography over silica gel eluted with petroleum ether- Me_2CO (1 : 0 \rightarrow 1 : 1) to give fraction Fr. 1–7. Fr. 3 (50 g) was repeatedly separated on silica gel, Rp-18, and Sephadex LH-20 (MeOH) column chromatography to afford **2** (19 mg), **3** (40 mg), and **4** (30 mg). Fr. 6 (45 g) was chromatographed on silica gel column eluting with CHCl_3 -MeOH (30 : 1 \rightarrow 1 : 1) and Rp-18 column eluting with a MeOH- H_2O gradient system (20% \rightarrow 60%), further purified on Sephadex LH-20 using MeOH as solvent to yield **1** (15 mg).

syringic acid-4-O- α -L-rhamnopyranoside (1): $\text{C}_{15}\text{H}_{19}\text{O}_9$, yellow amorphous powder. $[\alpha]_{\text{D}}^{26} = -81.20^\circ$ ($c = 0.12$, MeOH). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3422, 2936, 2851, 1700, 1592, 1553, 1502, 1460, 1393, 1129. UV (CH_3OH) λ_{max} ($\log \epsilon$): 211 (4.38), 254 (3.89) nm. ^1H -NMR and ^{13}C -NMR: see Table 1. Negative FAB-MS m/z (%): 343 $[\text{M}-1]^-$ (83), 325 $[\text{M}-18]^-$ (100), 197 $[\text{M}-\text{H}-146]^-$ (43); HR-ESI-MS m/z : 343.1040 $[\text{M}-1]^-$ (calcd for $\text{C}_{15}\text{H}_{18}\text{O}_9$ calcd. 343.1029).

1 β , 4 β -dihydroxy-5 α , 8 β (H) -eudesm-7 (11) Z-en-8, 12-olide (2): $\text{C}_{15}\text{H}_{22}\text{O}_4$, white powder. ^1H -NMR (500 MHz, CD_3OD): δ_{H} 4.94 (1H, m, H-8), 3.24 (1H, dd, $J = 3.8, 11.9$ Hz, H-1), 2.92 (1H, dd, $J = 3.5, 14.1$ Hz, H-6 α), 2.64 (1H, dd, $J = 6.3, 11.8$ Hz, H-9 β), 2.54 (1H, t, $J = 13.6$ Hz, H-6 β), 1.90 (1H, m, H-2 β), 1.81 (3H, m, H-13), 1.72

(1H, m, H-3 β), 1.55 (1H, m, H-2 α), 1.46 (1H, m, H-3 α), 1.23 (3H, s, H-15), 1.20 (3H, s, H-14), 1.16 (1H, dd, J = 3.5, 13.1 Hz, H-5), 0.89 (1H, t, J = 11.7 Hz, H-9 α); ^{13}C -NMR (125 MHz, CD_3OD): δ_{C} 79.1 (d, C-1), 27.2 (t, C-2), 40.1 (t, C-3), 71.6 (s, C-4), 52.6 (d, C-5), 24.0 (t, C-6), 166.4 (s, C-7), 80.4 (d, C-8), 47.2 (t, C-9), 41.5 (s, C-10), 120.3 (s, C-11), 177.3 (s, C-12), 8.0 (q, C-13), 13.4 (q, C-14), 29.7 (q, C-15).

1 β , 4 α -dihydroxy-5 α , 8 β (H)-eudesm-7 (11) Z-en-8, 12-olide (3): $\text{C}_{15}\text{H}_{22}\text{O}_4$, white powder. ^1H -NMR (500 MHz, CD_3OD): δ_{H} 4.66 (1H, dd, J = 6.3, 10.4 Hz, H-8), 3.15 (1H, dd, J = 4.3, 11.3 Hz, H-1), 2.96 (1H, dd, J = 3.3, 13.9 Hz, H-6 α), 2.55 (1H, dd, J = 6.2, 11.9 Hz, H-9 β), 2.12 (1H, m, H-6 β), 1.74 (1H, m, H-3 α), 1.68 (3H, s, H-13), 1.57 (2H, m, H-2), 1.48 (1H, m, H-3 α), 1.19 (1H, dd, J = 13.2, 3.4 Hz, H-5), 1.07 (3H, s, H-15), 0.91 (3H, s, H-14), 0.80 (1H, t, J = 11.7 Hz, H-9 α); ^{13}C -NMR (125 MHz, CD_3OD): δ_{C} 77.3 (d, C-1), 27.7 (t, C-2), 40.1 (t, C-3), 71.9 (s, C-4), 53.0 (d, C-5), 22.4 (t, C-6), 163.7 (s, C-7), 78.4 (d, C-8), 46.8 (t, C-9), 39.9 (s, C-10), 119.4 (s, C-11), 175.7 (s, C-12), 7.6 (q, C-13), 13.0 (q, C-14), 21.6 (q, C-15).

(-) - (7S, 8R) -dihydrodehydrodiconifery alcohol (4): $\text{C}_{20}\text{H}_{24}\text{O}_6$, colorless oil. ^1H -NMR (CD_3OD , 400 MHz): δ_{H} 6.94 (1H, d, J = 1.8, H-2), 6.82 (1H, dd, J = 1.8, 8.1 Hz, H-6), 6.78 (1H, s), 6.75 (1H, s, H-2'), 6.72 (1H, s, H-6'), 5.49 (1H, d, J = 6.3 Hz, H-7), 3.84 (3H, s, OCH_3), 3.80 (3H, s, OCH_3), 3.75 (1H, m, H-9), 3.56 (2H, t, J = 6.5 Hz, H-9'), 3.46 (1H, m, H-8), 2.62 (2H, t, J = 7.4 Hz, H-7'), 1.83 (2H, m, H-8'); ^{13}C -NMR (CD_3OD , 100 MHz): δ_{C} 134.8 (C-1), 110.5 (C-2), 147.5 (C-3), 145.2 (C-4), 116.1 (C-5), 119.7 (C-6), 89.0 (C-7), 55.4 (C-8), 64.9 (C-9), 136.9 (C-1'), 114.0 (C-2'), 145.2 (C-3'), 149.1 (C-4'), 129.8 (C-5'), 117.9 (C-6'), 35.8 (C-7'), 32.9 (C-8'), 62.2 (C-9'), 56.3, 56.7 (3, 3'- OCH_3).

Acknowledgements: The authors are grateful to the members of the analytical group of the State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences for all of the spectral measurements.

Reference:

- Editorial Committee of the Administration Bureau of Traditional Chinese Medicine. 1998, *Zhonghua Benchao* (中华本草) [M]. Shanghai: Shanghai Science and Technology Press, **3**: 453—454
- Jiangsu New Medical College. 1977. *Dictionary of Traditional Chinese Medicine* (中药大辞典) [M]. Shanghai: Shanghai Science and Technique Press, 652
- Kasai R, Okihara M, Asakawa J *et al.*, 1979. ^{13}C NMR study of α - and β -anomeric pairs of D-mannopyranosides and L-rhamnopyranosides [J]. *Tetrahedron*, **35** (11): 1427—1432
- Li CJ, Zhang DM, Luo YM *et al.*, 2008. Bis-sesquiterpenes and diterpenes from *Chloranthus henryi* [J]. *Phytochemistry*, **69** (16): 2867—2874
- Shao L (邵莉), Huang WH (黄卫华), Zhang CF (张朝凤) *et al.*, 2008. Study on chemical constituents from stem of *Dendrobium aphyllum* [J]. *China Journal of Chinese Materia Medical* (中国中药杂志), **33** (14): 1693—1695
- Teng F, Zhong HM, Chen CX *et al.*, 2009. Four new eudesmane sesquiterpenoid lactones from *Chloranthus serratus* [J]. *Helvetica Chimica Acta*, **92**: 1298—1303
- Wang XC, Zhang YN, Wang LL *et al.*, 2008. Lindenane sesquiterpene dimers from *Chloranthus fortunei* [J]. *Journal of Natural Products*, **71** (4): 674—677
- Wu B, He S, Wu X *et al.*, 2007. Cadinane and eudesmane sesquiterpenoids from *Chloranthus henryi*. *Helvetica Chimica Acta*, **90** (8): 1586—1592
- Xu YJ, Tang CP, Ke CQ *et al.*, 2007. Mono- and di-sesquiterpenoids from *Chloranthus spicatus* [J]. *Journal of Natural Products*, **70** (12): 1987—1990
- Yang SP, Yue JM, 2006. Chloramultilide A, a highly complex sesquiterpenoid dimer from *Chloranthus multistachys* [J]. *Tetrahedron Letters*, **47** (7): 1129—1132
- Yang SP, Zhang CR, Chen HD *et al.*, 2007. Sesquiterpenoids from *Chloranthus spicatus* (Thunb.) Makino [J]. *Chinese Journal of Chemistry*, **25** (12): 1892—1895
- Yang SP, Gao ZB, Wu Y *et al.*, 2008. Chlorahololides C-F: a new class of potent and selective potassium channel blockers from *Chloranthus holostegius* [J]. *Tetrahedron*, **64** (9): 2027—2034
- Yang SP, Yue JM, 2008. Terpenoids from *Chloranthus multistachys* [J]. *Natural Product Research*, **22** (13): 1163—1168
- Zhou ZK (周浙昆), 1993. Origin, systematics and distribution of Chloranthaceae [J]. *Acta Botanica Yunnanica* (云南植物研究), **15**: 321—331
- Zhu LP (朱丽萍), Li Y (李媛), Yang JZ (杨敬芝) *et al.*, 2008. Study on the chemical constituents of *Sarcandra glabra* [J]. *China Journal of Chinese Materia Medica* (中国中药杂志), **33** (2): 155—157